



# eclipse®

COMBINATION POUR-ON FOR CATTLE



Active Constituents: ABAMECTIN 10 mg/mL LEVAMISOLE 200 mg/mL

## Product description\*

For the treatment and control of roundworms, including macrocyclic lactone or levamisole resistant strains, and external parasites in cattle.

## Product benefits

- ✓ Recent field studies showed an average efficacy of over 99.5% against roundworms, including resistant strains, on Australian farms tested<sup>1</sup>
- ✓ The only pour-on with a registered claim to control resistant worms
- ✓ Convenient low 1 mL per 20 kg dose rate
- ✓ 2 actives, 1 pour-on
- ✓ Rainfast

## Broad-spectrum activity against resistance

ECLIPSE is Australia's first dual action broad-spectrum pour-on for cattle.

Being a dual action pour-on, the abamectin component works to kill a broad spectrum of internal and external parasites, while the levamisole component works to kill ML-resistant strains of *Cooperia* and *Ostertagia*.

By combining two powerful actives in one single convenient dose, you now have the power of a combination in a pour-on.

## Cattle Indications

For the treatment and control of adult and immature stages of\*:

### Gastrointestinal roundworms, including strains resistant to macrocyclic lactones ("mectin drenches") or to imidazothiazoles ("clear drenches" such as levamisole)

Barber's Pole Worm (*Haemonchus placei*)  
Small Brown Stomach Worm (*Ostertagia ostertagi*, *O. lyrata*, including inhibited stages)  
Stomach Hair Worm (*Trichostrongylus axei*)  
Small Intestinal Worm (*Cooperia oncophora*, *C. punctata*, *C. pectinata*, *C. mcmasteri*)  
Hookworm (*Bunostomum phlebotomum*)  
Thin-necked Intestinal Worm (*Nematodirus* spp.)  
Intestinal Threadworm (*Strongyloides papillosus*)  
Black Scour Worm (*Trichostrongylus* spp.)  
Large Bowel Worm (*Oesophagostomum radiatum*)  
Large Mouth Bowel Worm (*Chabertia ovina*)  
Whipworm (*Trichuris* spp.)

### Lungworms (*Dictyocaulus viviparus*)

### Lice

Long-nosed Sucking Lice (*Linognathus vituli*)  
Biting Lice (*Bovicola (Damalinia) bovis*)

### Buffalo Fly (*Haematobia irritans exigua*)

ECLIPSE Pour-On controls infection with Buffalo Fly for up to 10 days after treatment.

### Cattle Tick (*Boophilus microplus*)

Maximum tick efficacy is achieved 4-5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control program.



## Directions for use

### Dosage and Administration

ECLIPSE Pour-On dose rate is 1 mL / 20 kg body weight (0.5 mg/kg abamectin and 10 mg/kg levamisole). Apply along the backline of the animal, using the ECLIPSE Pour-On applicator, in a strip starting between the shoulder blades.

Only use the pour-on applicator supplied for use with ECLIPSE Pour-On. ECLIPSE is not compatible with other guns. After use, rinse gun with warm soapy water.

**Re-treatment interval:** DO NOT re-treat animals for 35 days after last treatment.

Liveweight (kg)	Dose Rate (mL)	1 Litre pack treats	2.5 Litre pack treats	5 Litre pack treats	10 Litre pack treats
51-100	0.5 mL / 10 kg	at least 200	at least 500	at least 1000	at least 2000
101-150	7.5	133	333	666	1333
151-200	10	100	250	500	1000
201-250	12.5	80	200	400	800
251-300	15	66	166	333	666
301-350	17.5	57	142	285	571
351-400	20	50	125	250	500
401-450	22.5	44	111	222	444
451-500	25	40	100	200	400
501-550	27.5	36	90	181	363
551-600	30	33	83	166	333
601-650	32.5	30	76	153	307

DO NOT USE in calves under 50 kg bodyweight. Under 100 kg dose at rate of 0.5 mL / 10 kg bodyweight, and over 650 kg dose at rate of 1 mL / 20 kg bodyweight.

## Withholding Periods

**MEAT:** DO NOT USE less than 56 days before slaughter for human consumption.

Calves MUST NOT be slaughtered for human consumption within 60 days of treatment of dams.

**MILK:** DO NOT USE in lactating cows or within 60 days of calving where milk may be used or processed for human consumption. If cows calve earlier than 60 days after treatment, milk may contain residues. This milk MUST NOT be used for human consumption, supplied for processing or fed to bobby calves for at least 60 days following treatment.

**EXPORT SLAUGHTER INTERVAL (ESI):** DO NOT USE less than 70 days before slaughter for export.

## Rainfast

ECLIPSE Pour-On is rainfast. Rain occurring as early as 2 hours after treatment is unlikely to affect the efficacy of the product. However rainfall on cattle less than 2 hours after treatment may result in reduced efficacy. ECLIPSE Pour-On should not be applied to wet animals.

## Storage

Store below 25°C (air conditioning).

Protect from light.

## Pack sizes

1L, 2.5L backpack, 5L backpack, 10L Gun Pack.

